

MEETING ABSTRACT

Open Access

Congruency of tumour volume delineated by FET PET and MRSI

Jörg Mauler^{1*}, Karl-Josef Langen¹, Andrew A Maudsley², Omid Nikoubashman³, Christian Filss¹, Gabriele Stoffels¹, N Jon Shah¹

From PSMR 2015: 4th Conference on PET/MR and SPECT/MR
La Biodola, Isola d'Elba, Italy. 17-21 May 2015

¹Institute of Neuroscience and Medicine, Forschungszentrum Jülich, Germany

In addition to MR imaging, PET imaging of O-(2-[¹⁸F]Fluorethyl)-L-Tyrosine (FET) uptake provides information on brain tumour extent and metabolic activity. Similarly, MRS has been shown to be of value for distinguishing high- from low-grade gliomas. Based on 2D spatially resolved MRSI, an overlap between ¹⁸FET uptake and the choline/N-acetyl-aspartate (Cho/NAA) ratio of more than 75 % has been reported.

Aim

To measure spatial correlation of ¹⁸FET-PET with 3D spatially resolved MRSI in patients with gliomas.

Methods

14 patients (46±16 y) with gliomas (WHO grade II-IV) were examined by simultaneous ¹⁸FET-PET-3D-MRSI measurements which covered the whole brain (Siemens Brain-PET/3T MR TIM Trio; MRSI: EPSI sequence, TE=17.6 ms). The data were analysed with respect to the congruency of the suspicious tissue delineated by the FET uptake and the Cho/NAA ratio. Locations of individual maxima and distances in between were determined. The congruency of the tumour was assessed using Dice's coefficients for assumed same tumour volume. All comparisons were carried out at the spatial resolution of the whole brain spectroscopic image (64×64×32 v×l, 5.6×5.6×10mm³ each).

Results

The intensity maxima of both modalities were (61±51) mm distant from each other. The average level of congruency between the tumour volumes delineated in the FET uptake and MRSI data was (33±25)%. Metabolically active tumour tissue, as depicted by FET uptake, is represented to a low extent by the choline/N-acetyl-aspartate ratio measured by spatially resolved 3D MRSI, which is in contrast to previous findings. Both modalities may reflect independent physiological properties of gliomas.

Authors' details

¹Institute of Neuroscience and Medicine, Forschungszentrum Jülich, Germany. ²Miller School of Medicine, University of Miami, USA. ³Department of Neuroradiology, Faculty of Medicine, RWTH Aachen University, Germany.

Published: 18 May 2015

doi:10.1186/2197-7364-2-S1-A61

Cite this article as: Mauler et al.: Congruency of tumour volume delineated by FET PET and MRSI. *EJNMMI Physics* 2015 **2**(Suppl 1):A61.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
